

A long-time limit for world subway networks

Camille Roth, Soong Moon Kang, Michael Batty and Marc Barthelemy

J. R. Soc. Interface published online 16 May 2012
doi: 10.1098/rsif.2012.0259

References

This article cites 19 articles, 2 of which can be accessed free

<http://rsif.royalsocietypublishing.org/content/early/2012/05/15/rsif.2012.0259.full.html#ref-list-1>

P<P

Published online 16 May 2012 in advance of the print journal.

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right-hand corner of the article or click [here](#)

Advance online articles have been peer reviewed and accepted for publication but have not yet appeared in the paper journal (edited, typeset versions may be posted when available prior to final publication). Advance online articles are citable and establish publication priority; they are indexed by PubMed from initial publication. Citations to Advance online articles must include the digital object identifier (DOIs) and date of initial publication.

To subscribe to *J. R. Soc. Interface* go to: <http://rsif.royalsocietypublishing.org/subscriptions>

A long-time limit for world subway networks

Camille Roth^{1,2,*}, Soong Moon Kang³, Michael Batty⁴
and Marc Barthelemy^{1,5}

¹*CAMS (CNRS/EHESS) 190, avenue de France, 75013 Paris, France*

²*Institut des Systèmes Complexes Paris-Ile de France (ISC-PIF), 57-59 rue Lhomond, 75005 Paris, France*

³*Department of Management Science and Innovation, University College London (UCL), Gower Street, London WC1E 6BT, UK*

⁴*Centre for Advanced Spatial Analysis (CASA), University College London, 90 Tottenham Court Road, London W1N 6TR, UK*

⁵*Institut de Physique Théorique, CEA, IPhT, CNRS-URA 2306, 91191 Gif-sur-Yvette, France*

We study the temporal evolution of the structure of the world's largest subway networks in an exploratory manner. We show that, remarkably, all these networks converge to a shape that shares similar generic features despite their geographical and economic differences. This limiting shape is made of a core with branches radiating from it. For most of these networks, the average degree of a node (station) within the core has a value of order 2.5 and the proportion of $k = 2$ nodes in the core is larger than 60 per cent. The number of branches scales roughly as the square root of the number of stations, the current proportion of branches represents about half of the total number of stations, and the average diameter of branches is about twice the average radial extension of the core. Spatial measures such as the number of stations at a given distance to the barycentre display a first regime which grows as r^2 followed by another regime with different exponents, and eventually saturates. These results—difficult to interpret in the framework of fractal geometry—confirm and yield a natural explanation in the geometric picture of this core and their branches: the first regime corresponds to a uniform core, while the second regime is controlled by the interstation spacing on branches. The apparent convergence towards a unique network shape in the temporal limit suggests the existence of dominant, universal mechanisms governing the evolution of these structures.

Keywords: evolution of networks; urban transportation; spatial networks; core and branch geometry

1. INTRODUCTION

Transportation systems, especially mass transit, are an important component in cities and their expansion. In a world where more than 50 per cent of the population lives in urban areas [1], and where individual transportation increases in cost as cities grow larger, mass transit and, in particular, subway networks are central to the evolution of cities, their spatial organization [2–4] and dynamical processes occurring in them [5,6]. The percentage $s(P)$ of cities with a subway system versus their population size P is shown in figure 1 (the data were obtained for cities with a population larger than 10^5 (<http://unstats.un.org/>)), which confirms that the larger a city, the more likely it is to have some form of mass transit system (see also [7]). Approximately 25 per cent of the cities having more than one million individuals, 50 per cent of those having more than two million and all those having

above 10 million have a subway system (as an indication, an exponential fit of the plot in figure 1 gives $s(P) = 1 - \exp(-P/P_0)$ where the typical population P_0 is in the order of three million).

For some cities, subway systems have existed for more than a century. Fascination with the apparent diversity of their structure has led to many studies and to particular abstractions of their representation in the design of idealized transit maps [8], and although these might appear to be planned in some centralized manner, it is our contention here that subway systems like many other features of city systems evolve and self-organize themselves as the product of a stream of rational but usually uncoordinated decisions taking place through time.

Generally speaking, subway systems have been developed to improve movement in urban areas and to reduce congestion. The early history of subways is sometimes connected to large-scale planning, for instance with the need to bring people from a growing periphery to the centre of the city where traditionally

*Author for correspondence (roth@ehess.fr).




Figure 1. Percentage of cities with a subway system versus the population (data from the UN (<http://unstats.un.org/>)). (Online version in colour.)

production and exchange usually take place. More broadly, it might seem that subway systems are engineered systems and intentionally structured in a core/periphery shape with their self-organization thus playing only a very minor role. This actually would be true if these subway systems were planned from their beginning to their current shape, but this is not the case for most networks. Their shape results from multiple actions, from planning within a time-limited horizon, set within the wider context of the evolution of the spatial distribution of population and related economic activities. We thus conjecture that subway networks actually result from a superimposition of many actions, both at a central level with planning and at a smaller scale with the reorganization and regeneration of economic activity and the growth of residential populations. In this perspective, subway systems are self-organizing systems, driven by the same mechanisms and responding to various geographical constraints and historical paths. This self-organized view leads to the idea that—beside local peculiarities due to the history and topography of the particular system—the topology of world subway networks display general, universal features within the limits of the physical geometry and cultural context in which their growth takes place.

The detection and characterization of these features require us to understand the evolution of these spatial structures. Indeed, subway networks are spatial [9–11] in the sense that they form a graph where stations are the nodes and links represent rail connections. We now understand quite well how to characterize a spatial network but we still lack tools for studying their temporal evolution. The present article tackles this problem, proposing various measures for these time-dependent, spatial networks.

Here we focus on the largest networks in major world cities and thus ignore currently developing, smaller networks in many medium-sized cities. We thus consider most of the largest metro networks (with at least one hundred stations) which exist in major world cities. These are: Barcelona, Beijing, Berlin, Chicago, London, Madrid, Mexico, Moscow, New York City

(NYC), Osaka, Paris, Seoul, Shanghai and Tokyo, for which we show a sample in figure 2. Additionally, we focus on urban subway systems and do not consider longer-distance heavy and light-rail commuting systems in urban areas, such as Réseau Express Régional (RER) in Paris or overground NetworkRail in London.

Static properties of transportation networks have been studied for many years [12] and, in particular, simple connectivity properties were studied in [13], while fractal aspects were considered in [14]. With the recent availability of new data, studies of transportation systems have accelerated [11] and this is particularly so for subway systems [15–23]. These studies have revealed some significant similarities between different networks, despite differences in their historical development and in the cultures and economies in which they have been developed. In particular, their average shortest path seems to scale with the square root of the number of stations and the average clustering coefficient is large, consistent with general results associated with two-dimensional spatial networks [11]. In the study of Sienkiewicz & Holyst [17], a strong correlation between the number of stations (for bus and tramway systems) and population size has been observed for 22 Polish cities, but such correlations are not observed at the world level (for all public transportation modes [21]).

Our empirical analysis of the evolution of these transportation networks is in line with approaches developed in the 1970s (see [24] and references therein) but we take advantage here of recent progress made in the understanding of spatial networks in general and new historical data sources that provide us with detailed chronologies of how these networks have developed.

1.1. Data

The network topologies at various points in time were built using two main data sources. First, current network maps as for 2009 were used to define lines for each network, and then to define line-based topologies, i.e. which station(s) follow(s) which other station(s) on each line. This information was then combined with opening dates for lines and stations. This second type of data has been gathered from Wikipedia (<http://www.wikipedia.org>): for most networks, there is one page per station with various information, including the first date of operation, the precise location and address, number of passengers, etc. The network building process for a given year t is then as follows. The list of open lines at year t is first established. For each open line, open stations at year t are listed and connections are created between contiguous stations according to the network topology. A station that is not open at year t on the given line, even if it is already open on a different line, is evidently discarded for the construction of the line. Eventually, those independent line topologies are gathered into the subway graph corresponding to year t . Note that we used 2009 topologies as it was relatively difficult to find and process network maps for all these networks for each year of their existence. As a result, topologies for any given year before 2009 may overlook topology features pertaining to station or line closures: for instance, a station that existed




Figure 2. A sample of large subway networks in large urban areas, all displaying a core and branches structure. From left to right and top to bottom: Shanghai, Madrid, Moscow, Tokyo, Seoul, Barcelona (figures from Wikimedia Commons (<http://commons.wikimedia.org>)). (Online version in colour.)

Table 1. List of various indicators (for the year 2009) for the major subway networks considered in this study (and sorted according to their metro population). P is the metropolitan area population (for 2009). N_L is the number of lines, N the number of physical stations, $\bar{\ell}_1$ is the average interstation distance, ℓ_T total route length, ℓ_T^{reg} the total route length for a regular graph with the same average degree, area and number of stations and β the final ratio between branch and core stations.

city	P (millions)	N_L	N	$\bar{\ell}_1$ (km)	ℓ_T (km)	$\ell_T/\ell_T^{\text{reg}}$	β (%)
Beijing	19.6	9	104	1.79	204	0.14	39
Tokyo	12.6	13	217	1.06	279	0.13	43
Seoul	10.5	9	392	1.39	609	0.39	38
Paris	9.6	16	299	0.57	205	0.18	38
Mexico City	8.8	11	147	1.04	170	0.15	39
NYC	8.4	24	433	0.78	373	0.12	36
Chicago	8.3	11	141	1.18	176	0.08	71
London	8.2	11	266	1.29	397	0.20	47
Shanghai	6.9	11	148	1.47	233	0.21	61
Moscow	5.5	12	134	1.67	260	0.16	71
Berlin	3.4	10	170	0.77	141	0.30	60
Madrid	3.2	13	209	0.90	215	0.42	46
Osaka	2.6	9	108	1.12	137	0.88	43
Barcelona	1.6	11	128	0.72	103	0.32	38

between 1900 and 1940 and that remained closed until now will not appear in any of our network datasets (such is the case for the British Museum Tube station). We suggest, however, that the effect of this bias is limited: on the one hand, generally few stations undergo closure in the course of the network evolution; on the other hand, these stations are rarely hubs, most often intermediary stations (of degree 2, i.e. connected to two stations), thus their non-inclusion bears little topological impact.

2. EXPLORING STATIC PROPERTIES

The main characteristics of the networks we have chosen are shown in table 1 where we first observe that the number of different lines appears to increase incrementally with the number of stations and that on average,

for these world networks, there are approximately 18 stations per line. Also, the mean interstation distance is on average $\bar{\ell}_1 \approx 1$ km with Beijing and Moscow showing the longest ones (1.79 and 1.67 km, respectively), and Paris displaying the shortest one (570 m), a diversity which finds its origin in the different historical paths of these networks. Other quantities such as the catchment area (the average number of individuals served by one station) could be computed but should be used with care: residential and economic activity density vary strongly across space and back-of-the-envelope arguments should only serve as a guide. Generally speaking, many parameters such as the population density, land use activity distribution and traffic are important drivers in the evolution of those networks, but we will focus in this first study on the characterization of these networks in terms of space and topology, independently of other

socio-economical considerations. A later extension of this research could examine these physical and topological properties with respect to various definitions of density which might include different activity types and various combinations related to the traffic that they generate.

To get some initial insight into the topology of these networks, one can first compare the total length ℓ_T of these networks to the corresponding quantity computed for an almost regular graph ℓ_T^{reg} with the same number of stations, area and average degree (the ‘degree’ of a node is the number of its neighbours in a graph). For a random planar graph with small degree fluctuations ($k \approx \langle k \rangle$) and small fluctuations of the spatial distribution of nodes, we can consider that the internode spacing is roughly constant and given by $\ell_0 \sim 1/\sqrt{\rho}$ where $\rho = N/A$ is the density of nodes defined as the number of nodes over the total area comprising all the nodes. The total length is then the number of edges $E = N\langle k \rangle/2$ times ℓ_0 which leads to [11]

$$\ell_T^{\text{reg}} \sim \frac{\langle k \rangle}{2} \sqrt{AN}. \quad (2.1)$$

In real applications, the determination of the quantity A is a difficult problem, but here we choose to use the metropolitan area as given by the various data sources. As shown in the table 1, the ratio $\ell_T/\ell_T^{\text{reg}}$ varies from 0.08 to 0.88, has an average of order 0.29 and displays essentially three outliers. First, Osaka (and also Madrid and Seoul) has a very large value indicating a highly reticulated structure. In contrast, Chicago and NYC have a much smaller value (≈ 0.1) signalling a more heterogeneous structure, which in both these cases is probably due to their strong geographical constraints.

The total length and the comparison with a regular structure gives a first hint about the structure of these networks but other indicators are needed to get a more focused view. There exist many different indicators and variables that describe these networks and their evolution. An important difficulty thus lies in the choice of the many possible indicators and how to extract useful information from them. In addition, the largest networks have a relatively small number of stations (always smaller than 500), which implies that we cannot expect to extract useful information from the probability distributions of various quantities as the results are too noisy. We thus have to compute more globally structured indicators which are, however, sensitive to the usually small temporal variations associated with these networks. In the following, we will focus on a certain number of these indicators, which we consider to be the most informative at this point.

Finally, we will focus in this study on purely spatial and topological properties: we will consider the evolution in space of these subway networks and we will not consider any other parameters that might be used to characterize urban growth. Our study is exploratory and thus a first step towards the integration of the most important factors into this research and despite its simplicity, in that we focus almost entirely on geometrical attributes, we consider that the evolution of the topology encodes many different factors

Table 2. t_0 is the initial year considered here for the different subways networks, \bar{v} is the average velocity (number of stations built per year), σ_v is the standard deviation of v and f is the fraction of years of inactivity (no stations built).

city	t_0	\bar{v}	σ_v	f (%)
Beijing	1971	3.3	7.74	79
Tokyo	1927	2.8	5.47	51
Seoul	1974	11.2	14.9	20
Paris	1900	2.6	5.1	60
Mexico City	1969	3.7	5.9	55
NYC	1878	3.3	8.3	68
Chicago	1901	1.9	6.24	71
London	1863	2.3	3.8	48
Shanghai	1995	14.9	20.2	31
Moscow	1936	1.7	1.9	43
Berlin	1901	1.6	3.3	65
Madrid	1919	2.3	4.6	59
Osaka	1934	1.4	4.1	79
Barcelona	1914	1.4	4.8	78

and that its study can point to some important general mechanisms governing the evolution of these networks.

3. NETWORK DYNAMICS

To get an initial impression of the dynamics of these networks, we first estimate the simplest indicator $v = dN/dt$, which represents the number of new stations built per year. From the instantaneous velocity, we can compute the average velocity over all years. This average can, however, be misleading as there are many years where no stations are built and thus we describe this by the fraction of ‘inactivity’ time f . We provide results for the networks considered in table 2 from which some interesting facts are revealed. Note that it is clear that Shanghai and Seoul are the most recent subway networks experiencing a rapid expansion that has elevated them to among the largest networks in the world.

For most of these networks, the average velocity is in a small range (typically $\bar{v} \in [1.4, 3.7]$) except for Seoul and Shanghai which are more recently developed networks. This is, however, an average velocity and we observe that (i) for all networks, larger velocities occur at earlier stages of the network and (ii) large fluctuations occur from one year to another. Interestingly, the fraction of the inactivity time (i.e. the time when no stations are built) is similar for all these networks with an average of about 58 per cent. We also show in figure 3a, the time evolution for each city of the number of stations, using an absolute timescale. In particular, the sizes of the oldest networks seem to progressively reach a plateau.

To make growth comparable across all networks, we introduce a second graph on figure 3b featuring the average, over all networks, of the number of stations after a certain number of years since the creation of the network. This average quantity exhibits a linear increase which indicates convincingly that, overall, as these networks become large, for a few decades thereafter, new stations represent an increasingly small percentage of the existing ones. In other words, the time evolution of all these networks is characterized




Figure 3. (a) Evolution of the number of stations for various large world subway networks. (b) Evolution of the number of stations y years after creation, averaged over all networks (tubes mark the standard deviation across all networks). The linear shape indicates that the growth in terms of new stations from a decade to another goes to zero for all these networks, signalling the possible appearance of a stationary limit. (Online version in colour.)

by small additions and not by sudden, abrupt changes with a large number of stations added in a small time duration. This first result anticipates the fact that these large networks may reach some kind of limiting shape that we will characterize in §3.1. This incremental growth of subways might reflect socio-economical concerns and pressure on the transportation networks such as diminishing return on investments as noted by various authors (see, [25] for US highways).

Finally, when we study the evolution of various indicators versus the number N of stations, an important point for our statistical analysis will be the number of subways with a given number N at a given time t . We show this quantity in figure 4 and we can see that, for $N \in [25, 100]$ approximately, this number is the largest (almost 15—note that this figure is nonetheless too small to allow a discussion of the normality of the various quantities considered below). Unfortunately, for larger values of N the number of cities is naturally smaller, and at this stage we cannot give definitive answers but suggest some limits for large N .

3.1. Characterization of the core and branches structure

The large subway networks considered here thus converge to a long-time limit where there is always an increasingly smaller percentage of new stations added through time. The remarkable point that we will show below is that all these networks, despite their geographical and economical differences, converge to a shape that exhibits several typical topological and spatial features. Indeed, by inspection, we observe that in most large urban areas, the network consists of a set of stations delimited by a ‘ring’ that constitute the ‘core’. From this core, quasi-one-dimensional branches grow and reach out to areas of the city further and further from the core. In figure 2, we show a sample of these networks as they currently exist. We note here that the ring, which is defined topologically as the set of core stations that are either at the junction of branches or on the shortest geodesic path connecting these junction stations, exists or not as a subway line. For




Figure 4. Number of cities with a given number of stations at a given time.

instance, for Tokyo, there is a such a circular line (called the Yamanote line), while for Paris the topological ring does not correspond to a single line. It is also worth noting that in those systems where the core is harder to define such as NYC where physical constraints are strongly manifest (the east and west rivers which bound Manhattan), a pseudo core is evident where a series of lines coalesce to enable travellers to move around the core circumferentially.

More formally, branches are defined as the set of stations that are iteratively built from a ‘tail’ station, or a station of degree 1. New neighbours are added to a given branch as long as their degree is 2—continuing the line, or 3—defining a fork. In this latter case, the aggregative process continues *if and only if* at least one of the two possible new paths stemming from the fork is made up of stations of degree 2 or less. Note that the core of a network with no such fork is thus a k -core with $k = 2$ [26].

The general structure can schematically be represented as in figure 5.

We first characterize this branch and core structure with the parameter $\beta(t)$ defined as

$$\beta(t) = \frac{N_B(t)}{N_B(t) + N_C(t)}, \quad (3.1)$$




Figure 5. Schematic of subway networks. A large ‘ring’ encircles a core of stations. Branches radiate from the core and reach further areas of the urban system. The branches are essentially characterized by their size (parameter $\beta(t)$, equation (3.1)), and their spatial extension (parameter $\eta(t)$ in equation (3.2)). The core is characterized by its average degree $\langle k_{\text{core}} \rangle(t)$ defined in equation (3.3) and fraction of nodes of degree 2 (f_2), its number of stations $N_C(t)$ and its size $r_C(t)$. Green circles denote junction; red, terminus. (Online version in colour.)

where $N_B(t)$ and $N_C(t)$, respectively, represent the number of stations on branches and the number of stations in the core at time t .

We can also further characterize the structure of branches. Their topological properties are trivial and their complexity resides in their spatial structure. We can then determine the average distance (in kilometres) from the geographical barycentre of the city to all core and branch stations, respectively: $\bar{D}_C(t)$ and $\bar{D}_B(t)$ (the barycentre is computed as the centre of mass of all stations, or in other words, the average location of all the stations). This last distance provides information about the spatial extension of the branches when we can form the ratio $\eta(t)$:

$$\eta(t) = \frac{\bar{D}_B(t)}{\bar{D}_C(t)} \quad (3.2)$$

which gives a spatial measure of the amount of extension of the branches.

We also need information on the structure of the core. The core is a planar (which is correct at a good accuracy for most networks) spatial network and can be characterized by many parameters [11]. It is important to choose those that are not simply related but ideally represent different aspects of the network (such as those proposed in the form of various indicators; see, for example, [11,12,27]). At each time step t , we will characterize the core structure by the following two parameters. The first parameter is simply the average degree of the

core which characterizes its ‘density’:

$$\langle k_{\text{core}} \rangle(t) = \frac{2E_C(t)}{N_C(t)}, \quad (3.3)$$

where $N_C(t)$ is the number of core nodes and $E_C(t)$ the number of its edges. The average degree is connected to the standard index $\gamma(t) = E_C(t)/(3N_C(t) - 6)$ where the denominator is the maximum number of links admissible for a planar network [12].

The average degree of the core contains useful information about it, and there are many other quantities (such as standard indices such as α , etc.; see, [12]) which can give additional information. We will use another simple quantity which describes in more detail the level of interconnections in the core and which is given by the fraction f_2 of nodes in the core with $k=2$. In the case of the well-interconnected system, this fraction will tend to be small, while sparse cores with a few interconnections will have a larger fraction of $k=2$ nodes.

Once we know this fraction f_2 of $k=2$ nodes in the core, which characterizes the level of interconnection and the parameter $\eta(t)$, which characterizes the relative spatial extension of branches, we have key information on the intertwining of both topological and geographical features in such ‘core/branch’ networks.

3.2. Time evolution of β , $\langle k_{\text{core}} \rangle$, f_2 and η

The historical development of these networks is very different from one city to another and representing the evolution of a specific quantity versus time would probably not be particularly meaningful. Similarly, city networks often experience significant development in some particular years, while they experience little or no evolution for the rest of the time. To be able to compare the networks across time periods and cities, we propose to study their evolution in terms of the number of stations N that are constructed.

We first plot in figure 6*a* the parameter β as a function of N for the networks studied here. It is difficult to draw strong conclusions from this plot, but we can bin these data and represent the average value of β per bin and its dispersion as well (figure 6*b*). In this figure, we may see that the average value of β seems to stabilize slowly to some value in [0.35, 0.55].

It is also important to characterize the spatial importance of the branches. The parameter η gives a precious indication about their extension and we show in figure 7 the evolution of this parameter with N (the data are binned). This figure shows that in the interval where we have the largest number of subways, the average value of η is around 2 with relatively large fluctuations which seem to decrease with N .

The parameters β and η give an indication of the importance of the core but do not say anything about its structure. A first structural indication may be given by its average degree $\langle k_{\text{core}} \rangle$ and by the percentage of nodes in the core having a degree k equal to 2. In particular, these two quantities shed light on how interconnections are created in the core. We display in figure 8*a* the average degree of the core $\langle k_{\text{core}} \rangle$ which,




Figure 6. (a) Parameter β as a function of the number of stations N for the different world subways. (b) Same as (a) but averaged over 20 bins and showing the standard deviation. (Online version in colour.)




Figure 7. Evolution of the ratio η , which characterizes the spatial extension of branches relative to the core. (Online version in colour.)

even if there is a slow increase with N , displays moderate variations around 2.4 approximately.

This value is relatively small and indicates that the fraction of connecting stations (i.e. with $k > 2$) is also small and means that most core stations belong to one single line with few that actually allow connections. More precisely, we observe in figure 8b that, on average, for subways with $N < 100$ the fraction of interconnecting stations is increasing with N —which probably corresponds to some organization of the subway—but that for larger subways ($N > 100$), the percentage f_2 is increasing again, which probably corresponds to a densification process without the creation of new




Figure 8. (a) Average degree of the core $\langle k_{\text{core}} \rangle$ (equation (3.3)) and its dispersion versus number of stations (averaged over 20 bins). (b) Evolution of the percentage f_2 of $k=2$ core nodes (averaged over 20 bins). (Online version in colour.)




Figure 9. Evolution of the mean distance to the barycentre (in kilometres) of core stations with the number of stations N . (Online version in colour.)

interconnections. This densification can indeed be confirmed as the diameter of the core (figure 9) seems to reach a plateau for most cities.

As noted above, the number of subways with large N is smaller and the statistics therefore less reliable. At this point and with this statistical error in mind, we observe that the average value β and its dispersion are decreasing with N and suggests that β could converge to some ‘limiting’ value $\beta_{\infty} \approx 45$ per cent. The same remarks also apply to η and suggest a limiting value of order 2. Concerning the core, the dispersion of $\langle k_{\text{core}} \rangle$ is always moderate and approximately constant, showing that the fluctuations among different networks




Figure 10. The log–log plot of the number of different branches versus the number of stations for the different subway networks considered here. The dashed line is a power law fit with exponent ≈ 0.6 . (Online version in colour.)

are also moderate. We observe a slow increase of $\langle k_{\text{core}} \rangle$ pointing to a mild yet continuing densification of the core, even after a long period of time. The fraction of connecting stations has more complex dynamics and seems to decrease with N for large networks. In these networks, there is an obvious cost associated with the large value of $k > 2$ and such a decreasing fraction could be due to the fact that a small fraction is enough to enable easy navigation in the network.

In summary, our results display non-negligible fluctuations but suggest that large subway networks may converge to a long-time limiting network largely independent of their historical and geographical differences. So far, we can characterize the ‘shape’ of this long-time limiting network with values of $\beta_{\infty} \approx 45$ per cent, $\eta_{\infty} \approx 2$, and a core made of approximately 80 per cent of non-connecting stations. It will be interesting to observe the future evolution of these networks in order to confirm (or not) our current results.

3.3. Number of branches

We now consider the number \mathcal{N}_B of different branches. A naive argument would be that the number of branches is actually proportional to the perimeter of the core structure. This implicitly assumes that the distance between different branches is constant. In turn, the perimeter should roughly scale as \sqrt{N} as the core is a relatively dense planar graph and contains a number of nodes proportional to N . These assumptions thus lead to

$$\mathcal{N}_B \sim \sqrt{N} \quad (3.4)$$

We display the number of branches versus the number of stations N for the various networks considered here. A power law fit of the data presented in figure 10 gives $\mathcal{N}_B \sim N^b$ with $b \approx 0.6$ ($r^2 = 0.85$) consistent with our argument.

3.4. Balance between the core density and the branch structure

Even if it seems that the values of various indicators converge with the size of the networks, we still have




Figure 11. Relation between the spatial extension of branches and the degree of interconnection in the core. The 2009 values for the percentage f_2 of $k = 2$ core nodes and η are plotted for 12 city subways. (Online version in colour.)

appreciable variations. For example, η varies from ≈ 1 to ≈ 3 and exhibits a relatively constant and not negligible relative dispersion. It is thus important to understand the remaining differences between these networks. To achieve this, we focus on the relation between η which characterizes the spatial extension of the branches relative to the core, and the percentage f_2 of $k = 2$ nodes in the core which indicates how well connected the core is. We focus on the ‘final’ values of these parameters obtained for 2009 for the various networks and we obtain the plot shown in figure 11. From this figure, we first see that (η, f_2) ranges from $(\approx 1.4, \approx 85\%)$ for NYC up to $(\approx 3.3, \approx 45\%)$ for Moscow, which is indeed a highly ramified network with a very dense core.

Very roughly speaking, we first observe that, for this set of the largest subway systems in the world, the percentage f_2 is large and above 60 per cent and relatively independent from η . At a finer level, we observe from this figure that clusters of networks with similar properties also emerge. The first cluster comprises Beijing, Berlin, Shanghai and Seoul which are remarkably close to each other: f_2 is of order $80 \pm 5\%$ and $\eta(t) \approx 2.84 \pm 0.1$. This cluster corresponds thus to subway networks with a large degree of ramification and a lower interconnection level in their core. Not surprisingly, this cluster comprises rapidly evolving networks such as Beijing and Shanghai, for example. Another cluster comprises London, Paris and Madrid with a smaller value of $f_2 \approx 70 \pm 5\%$, which might result from their denser city centre structure and a smaller value of $\eta \approx 2$. This other cluster corresponds to denser networks, less ramified but with more interconnections in the core. Finally, we can identify another cluster made of Chicago and Osaka with a small value of η and a relatively dense core (with $f_2 \approx 70\%$).

4. SPATIAL ORGANIZATION OF THE CORE AND BRANCHES

Following earlier studies on the fractal aspects of subway networks [14], we can inspect the spatial subway organization by considering the number of stations $N(r)$ at a

Table 3. For each city, we compute the number of stations in the core (for the year 2009) and from the numerical calculation of $N(r)$ we can estimate r_C the size of the core (in kilometres) from $N(r = r_C) = N_C$.

City	N_C	r_C (km)
Beijing	63	4.4
Tokyo	123	5.0
Seoul	243	11.6
Mexico	90	4.7
Shanghai	57	3.7
Moscow	39	5.9
London	142	7.3
Paris	186	4.2
Madrid	113	4.4
Berlin	68	5.5
Barcelona	57	3.5
Osaka	46	3.6

distance less than or equal to r , where the origin of distances is the barycentre of all stations considered as points. Interestingly, the barycentre of all stations is almost motionless, except in the case of NYC where the barycentre moves from Manhattan to Queens, and thus we will exclude NYC from further study. Chicago is a similar case: the spatial structure of the core is peculiar, mainly due to the presence of the lake which constrains the network from expanding in the other directions. We will also exclude this network in this section. It should, however, be noted here that both Chicago and NYC do follow the image of core and branches but that the main difference with the other networks is that the core of these networks has no clear spatial meaning due to the geographical constraints (such as the presence of a lake for Chicago and a particular shape of the land area for NYC).

For the year 2009, the limiting shape made of a core and branches implies that there is an average distance r_C which determines the core. In practice, we can measure on the network the size N_C of the core and we then define r_C such that $N(r = r_C) = N_C$ (which assumes implicitly an isotropic core shape, which is the case for most networks except for the excluded cases of Chicago and NYC). For the various cities, we can easily compute the function $N = N(r)$ from which we can extract r_C and we report the results in table 3.

Next, we can rescale r by r_C and $N(r)$ by N_C and we then obtain the results shown in figure 12.

This figure displays several interesting features. First, the short distance regime $r < r_C$ is well described by a behaviour of the form $N(r) \sim \rho_C \pi r^2$ consistent with a uniform density ρ_C of core stations. For very large distances, for most networks we observe a saturation of $N(r)$. The interesting regime is then for intermediate distances when r is larger than the core size but smaller than the maximum branch size r_{\max} . This intermediate regime is characterized by different behaviours with r . A similar result was obtained earlier [14], where the authors observed for Paris that $N(r > r_C) \sim r^{0.5}$, a result that was at that time difficult to understand in the framework of fractal geometry.

Here we show that these regimes can be understood in terms of the core and branches model, with the




Figure 12. (a) Rescaled number of stations at distance r from the barycentre as a function of the rescaled variable r/r_C where r_C is the size of the core defined as $N(r = r_C) = N_C$ (shown here in log–log). The dotted line represents a power law $\sim r^2$ and serves as a guide to the eye. (b) The case of Moscow where the two regimes ($r < r_C$ and $r > r_C$) with their different exponents are visible (the dotted lines serve here as a guide to the eye). (Online version in colour.)

additional factor that the spacing between consecutive stations is increasing with r . Within this picture (and assuming isotropy), $N(r)$ is given by

$$N(r) \sim \begin{cases} \rho_C \pi r^2 & \text{for } r < r_C, \\ \rho_C \pi r_C^2 + \mathcal{N}_B \int_{r_C}^r \frac{dr}{\Delta(r)} & \text{for } r_C < r < r_{\max}, \\ N & \text{for } r > r_{\max}, \end{cases} \quad (4.1)$$

where N is the total number of stations, \mathcal{N}_B is the number of branches and $\Delta(r)$ is the average spacing between stations on branches at distance r from the barycentre.

To test this shape, we can determine the various parameters of equation (4.1)—namely \mathcal{N}_B , N_C , r_C and $\Delta(r)$ —and plot the resulting shape of equation (4.1) against the empirical data. It is easy to determine empirically the numbers \mathcal{N}_B , N_C and r_C , but the quantity $\Delta(r)$ is extremely noisy due to the small number of points (all these numbers are determined for the year 2009), especially for large values of r closest to r_{\max} , at a distance where, often, there is no more than a handful of stations.

The less noisy situation is obtained in the case of Moscow, which has long branches and for which we obtain a roughly constant interstation spacing. In this




Figure 13. $N(r)/N_C$ versus r/r_C for (a) Moscow, (b) Tokyo, (c) Paris and (d) Madrid (from top to bottom and left to right). The circles represent the data and the green solid line the fit using equation (4.1) with parameters estimated from the empirical data. (Online version in colour.)

case, we obtain, for $r > r_C$, a behaviour of the form $N(r) \sim \mathcal{N}_B r$ (figure 12*b*).

More generally, the large distance behaviour $r_C < r < r_{\max}$ will be of the form

$$N(r_C < r < r_{\max}) \sim r^{1-\tau} \quad (4.2)$$

where τ denotes the exponent governing the interspacing decay $\Delta(r) \sim r\tau$. For most networks, the regime $r_C < r < r_{\max}$ is small and as already mentioned $\Delta(r)$ is very noisy. Rough fits in different cases give a behaviour for equation (4.1) consistent with the data (figure 13).

In particular, for Moscow which has long branches, we observe a behaviour consistent with $\Delta(r) \simeq \text{constant}$, while for the other networks we observe an increasing trend, but an accurate estimate of τ is difficult to obtain, given the small variation range of r — with no more than one decade of available data. For example, a fit over this decade of data gives $\tau \approx 0.5$ for Paris (with $r^2 = 0.74$) in agreement with the result obtained in [14]. Despite the difficulty of obtaining accurate quantitative results, more data are needed to have a definite answer and so far we can only claim that the data are not inconsistent with the behaviour equation (4.1), which supports our picture of a long-time limit network shape made of a core and radial branches.

5. DISCUSSION

In summary, we have observed a number of similarities between different subway systems for the world's largest cities, despite their geographical and historical differences.

First, we have shown that the largest subway networks exhibit a similar temporal decrease of most fluctuations around their long-term stable values and

thus converge to a similar structure. We identified and characterized the shape of this long-time limiting graph as a structure made of a core and branches which appears to be relatively independent of the peculiar historical idiosyncracies associated with the evolution of these particular cities.

For large networks, we generally observe a fraction of branches of about 45 per cent for most networks, and a ratio for the spatial extensions of branches to the core of about 2. The number of branches scales roughly as the square root of the number of stations. The core of these different city networks has approximately the same average degree which is increasing with the network size, from ≈ 2 to ≈ 2.4 when $N \approx 100$, after which it approximately remains within the interval [2.3, 2.5] (with moderate fluctuations). The fraction of $k=2$ nodes in the core is generally larger than 60 per cent.

In addition, this picture of a core with branches and an increasing spacing between consecutive stations on these branches is confirmed by spatial measurements such as the number of stations at a given distance r and provides a natural interpretation to these measures.

The evolution of networks in general and urban networks in particular represents an exciting unexplored problem that mixes spatial and topological properties in unusual and often counterintuitive ways. They require a specific set of indicators that describe these phenomena. Other data such as population density, land use activity distribution and traffic flows are likely to bring relevant information to this problem and would undoubtedly enrich our study. We believe, however, that the present approach represents an important exploratory step in our understanding and is crucial for the modelling of the evolution of urban networks. In particular, the existence of unique long-time limit

topological and spatial features is a universal signature that fundamental mechanisms, independent of historical and geographical differences, contribute to the evolution of these transportation networks.

We thank the anonymous referees for very useful and interesting comments.

REFERENCES

- 1 Department of Economic and Social Affairs. 2011 UN Population division. <http://www.unpopulation.org>.
- 2 Hanson, S. & Giuliano, G. (eds) 2004 *The geography of urban transportation*. New York, NY: Guilford Publications.
- 3 Batty, M. 2005 *Cities and complexity*. Cambridge, MA: The MIT Press.
- 4 Niedzielski, M. A. & Malecki, E. J. In press. Making tracks: rail networks in world cities. *Ann. Assoc. Am. Geogr.* (doi:10.1080/00045608.2011.601212)
- 5 Bettencourt, L. M. A., Lobo, J., Helbing, D., Kuehnert, C. & West, G. B. 2007 Growth, innovation, scaling, and the pace of life in cities. *Proc. Natl Acad. Sci. USA* **104**, 7301–7306. (doi:10.1073/pnas.0610172104)
- 6 Balcan, D., Colizza, V., Gonçalves, B., Hu, H., Ramasco, J. & Vespignani, A. 2009 Multiscale mobility networks and the spatial spreading of infectious diseases. *Proc. Natl Acad. Sci. USA* **106**, 21 484–21 489. (doi:10.1073/pnas.0906910106)
- 7 Daganzo, C. F. 2009 Structure of competitive transit networks. *Transport. Res. B: Meth.* **44**, 434–446.
- 8 Ovenden, M. 2007 *Transit maps of the world*, 2nd edn and revised edition. London, UK: Penguin Books.
- 9 Reggiani, A. & Nijkamp, P. (eds) 2006 *Spatial dynamics, networks and modelling*. Cheltenham, UK: E. Elgar Publishing.
- 10 Reggiani, A. & Nijkamp, P. (eds) 2009 *Complexity and spatial networks: in search of simplicity*. Berlin: Springer.
- 11 Barthelemy, M. 2011 Spatial networks. *Phys. Rep.* **499**, 1–101. (doi:10.1016/j.physrep.2010.11.002)
- 12 Haggett, P. & Chorley, R. J. 1969 *Network analysis in geography*. London, UK: Edward Arnold.
- 13 Bon, R. 1979 Allometry in topologic structure of transportation networks. *Qual. Quan.* **13**, 307–326. (doi:10.1007/BF00188018)
- 14 Benguigui, L. & Daoud, M. 1991 Is the suburban railway system a fractal? *Geogr. Anal.* **23**, 362–369. (doi:10.1111/j.1538-4632.1991.tb00245.x)
- 15 Latora, V. & Marchiori, M. 2001 Is the Boston subway a small-world network? *Physica A* **314**, 109–113. (doi:10.1016/S0378-4371(02)01089-0)
- 16 Seaton, K. A. & Hackett, L. M. 2004 Stations, trains and small-world networks. *Physica A* **339**, 635–644. (doi:10.1016/j.physa.2004.03.019)
- 17 Sienkiewicz, J. & Holyst, J. A. 2005 Statistical analysis of 22 public transport networks in Poland. *Phys. Rev. E* **72**, 046127. (doi:10.1103/PhysRevE.72.046127)
- 18 Gattuso, D. & Miriello, E. 2005 Compared analysis of metro networks supported by graph theory. *Netw. Spat. Econ.* **5**, 395–414. (doi:10.1007/s11067-005-6210-5)
- 19 Angeloudis, P. & Fisk, D. 2006 Large subway systems as complex networks. *Physica A* **367**, 553–558. (doi:10.1016/j.physa.2005.11.007)
- 20 Lee, K., Jung, W.-S., Park, J. S. & Choi, M. Y. 2008 Statistical analysis of the Metropolitan Seoul Subway System: network structure and passenger flows. *Physica A* **387**, 6231–6234. (doi:10.1016/j.physa.2008.06.035)
- 21 von Ferber, C., Holovatch, T., Holovach, Y. & Palchykov, V. 2009 Public transport networks: empirical analysis and modeling. *Eur. Phys. J. B* **68**, 261–275. (doi:10.1140/epjb/e2009-00090-x)
- 22 Derrible, S. & Kennedy, C. 2010 The complexity and robustness of metro networks. *Physica A* **389**, 3678–3691. (doi:10.1016/j.physa.2010.04.008)
- 23 Derrible, S. & Kennedy, C. 2010 Characterizing metro networks: state, form, and structure. *Transportation* **37**, 275–297. (doi:10.1007/s11116-009-9227-7)
- 24 Xie, F. & Levinson, D. 2009 Topological evolution of surface transportation networks. *Comp. Environ. Urban Syst.* **33**, 211–223. (doi:10.1016/j.compenvurbysys.2008.09.009)
- 25 Naridi, I. & Manumeas, T. P. 1996 Contribution of highway capital to industrial and national productivity growth. Technical report. Work Order Number BAT-94-008. Federal Highway Administration (Office of Policy Development). See <http://www.fhwa.dot.gov/policy/gro98civr.htm>.
- 26 Seidman, S. B. 1983 Network structure and minimum degree. *Social Netw.* **5**, 269–287. (doi:10.1016/0378-8733(83)90028-X)
- 27 Xie, F. & Levinson, D. 2007 Measuring the structure of road networks. *Geogr. Anal.* **39**, 336–356. (doi:10.1111/j.1538-4632.2007.00707.x)